

IPL offers long lasting improvement in dry eye syndrome to complement ongoing lubricants and eyelid hygiene.

**Contraindications:** Please check if you have one of the following conditions:

- |  |  |
|--|--|
| <input type="checkbox"/> Skinphototype VI (very dark/black skin)                         | <input type="checkbox"/> Chronic or cutaneous viral, fungal, or bacterial diseases   |
| <input type="checkbox"/> Pregnancy   | <input type="checkbox"/> Anti-coagulant use  |
| <input type="checkbox"/> Heart Disease   | <input type="checkbox"/> Roaccutane  |
| <input type="checkbox"/> Epilepsy  | <input type="checkbox"/> Photosensitive medications  |
| <input type="checkbox"/> Active tans   | <input type="checkbox"/> Pacemaker   |
| <input type="checkbox"/> Keloid scarring under the eyelid                                | <input type="checkbox"/> Lupus disease   |
| <input type="checkbox"/> Porphyria   | <input type="checkbox"/> Gold Injections   |
| <input type="checkbox"/> Xeroderma   | <input type="checkbox"/> HIV/AIDS  |
| <input type="checkbox"/> Moles, birthmark or tattoos under the eyelid                    | <input type="checkbox"/> Haemophilia   |
| <input type="checkbox"/> Active Infections   | <input type="checkbox"/> Diabetes (note: treatment can be performed following a successful test patch monitored during 48 hours) |
| <input type="checkbox"/> Significant skin conditions or any inflammatory skin conditions | <input type="checkbox"/> Dysplastic Nevi   |
| <input type="checkbox"/> Damage to natural skin texture and/or very dry skin             | <input type="checkbox"/> Active cold sores, open lacerations or abrasions  |
|  | <input type="checkbox"/> Highly vasculated area in the lesion's immediate proximity  |

Please advise if you are taking any of these photo sensitive medications:

- |   |  |
|---|--|
| <input type="checkbox"/> St. Johns Wort medications   | <input type="checkbox"/> Declomycin chlortetracycline (Aureomycin)         |
| <input type="checkbox"/> Antibacterial group: sulphathiazole, long acting sulphonamides                   | <input type="checkbox"/> Chlorpromazine, promazine, trimeprazine, mepazine |
| <input type="checkbox"/> Diuretics: chlorothiazide, hydrochlorothiazide, quinethazone                     | <input type="checkbox"/> Psychotropic drugs                                |
| <input type="checkbox"/> Antidiabetic: sulphonylureas, cabamides  | <input type="checkbox"/> Chloroquine                                       |
| <input type="checkbox"/> Tetracyclines: tetracycline, doxycycline, dimethylchlortetracycline (Ledermycin) | <input type="checkbox"/> Antimalarials                                     |
| <input type="checkbox"/> Sulphonamides and related chemicals  | <input type="checkbox"/> Griseofulvin                                      |
| <input type="checkbox"/> Tomezathine hydrochloride  | <input type="checkbox"/> Amiodarone  |
| <input type="checkbox"/> Phenothiazines   | <input type="checkbox"/> Chlordiazepoxide                                  |
| <input type="checkbox"/> Diphenhydramine  | <input type="checkbox"/> Retinols, glycolic products, creams or oils       |